







Minimise future risk of exacerbations

Reduce fixed airflow limitation

Control side-effects of treatment

asthma patients have a x6 greater chance of having an asthma attack in the next few weeks than those with minimal to no daytime symptoms⁷

Symptomatic

symptom free days has been found to be the most important aspect for people with asthma8

Having



1. Demoly P. et al Eur Respir Rev 2010. 19. 116 150-157 2. Sastre J. et al. World Allergy Organisation Journal. 2016; 9:13 3. Price D et al. npj Primary Care Respiratory Medicine. 2014; 14009

4. Asthma UK. https://www.asthma.org.uk/about/media/facts-and-statistics/ (Accessed 26/07/2016) 5. Gruffyd-Jones K. et al. npj Primary Care Respiratory Medicine. 2016;16022 6. Selroos O et al. European Respiratory Review 2015 24: 474-483

7. Bateman ED, et al. J Allergy Clin Immunol. 2010 Mar;125(3):600-8 8. Lloyd A, et al. Prim Care Respir J. 2007;16:241-248

9. Global Initiative for Asthma (GINA). Global strategy for asthma management and prevention.

References:

Available from: http://www.ginasthma.org/local/uploads/files/GINA_Report_2014_Jun11.pdf (Accessed 30/07/16)